

MIT Joint Program on the Science and Policy of Global Change



Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content

A.P. Sokolov, C.E. Forest, and P.H. Stone

**Report No. 166
November 2008**

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.

Henry D. Jacoby and Ronald G. Prinn,
Program Co-Directors

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change
77 Massachusetts Avenue
MIT E40-428
Cambridge MA 02139-4307 (USA)

Location: One Amherst Street, Cambridge
Building E40, Room 428
Massachusetts Institute of Technology

Access: Phone: (617) 253-7492
Fax: (617) 253-9845
E-mail: globalchange@mit.edu
Web site: <http://mit.edu/globalchange/>

 Printed on recycled paper

Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content

A.P. Sokolov^{*}, C.E. Forest and P.H. Stone

Abstract

The MIT 2D climate model is used to make probabilistic projections for changes in global mean surface temperature and for thermosteric sea level rise under a variety of forcing scenarios. The uncertainties in climate sensitivity and rate of heat uptake by the deep ocean are quantified by using the probability distributions derived from observed 20th century temperature changes. The impact on climate change projections of using the smallest and largest estimates of 20th century deep ocean warming is explored. The impact is large in the case of global mean thermosteric sea level rise. In the MIT reference (“business as usual”) scenario the median rise by 2100 is 27 and 43 cm in the respective cases. The impact on increases in global mean surface air temperature is more modest, 4.9 C and 3.9 C in the two respective cases, because of the correlation between climate sensitivity and ocean heat uptake required by 20th century surface and upper air temperature changes. The results are also compared with the projections made by the IPCC AR4’s multi-model ensemble for several of the SRES scenarios. The multi-model projections are more consistent with the MIT projections based on the largest estimate of ocean warming. However the range for the rate of heat uptake by the ocean suggested by the lowest estimate of ocean warming is more consistent with the range suggested by the 20th century changes in surface and upper air temperatures, combined with expert prior for climate sensitivity.

Contents

1. INTRODUCTION	1
2. MODEL DESCRIPTION	2
3. PROBABILITY DISTRIBUTIONS OF INPUT CLIMATE PARAMETERS	4
4. SIMULATIONS	6
5. RESULTS AND DISCUSSION	8
6. CONCLUSIONS	12
7. REFERENCES	13

1. INTRODUCTION

There are significant uncertainties in the characteristics of the climate system which define its response to external forcing, such as climate sensitivity, strength of aerosol forcing and the rate of deep ocean heat uptake (*e.g.* Andronova and Schlesinger 2001; Frame *et al.* 2005; Forest *et al.* 2006; Knutti *et al.* 2006). Consequently anthropogenic climate change can be described only in probabilistic terms, even when changes in the concentrations of greenhouse gases (GHGs) and aerosols are prescribed (*e.g.* Knutti *et al.* 2003). To account for such uncertainties, projections of future changes in global mean surface air temperature (SAT) presented in the IPCC AR4 (Meehl *et al.* 2007) are based in part on a multi-model ensemble of simulations with coupled atmosphere-ocean general circulation models (AOGCMs).

^{*} MIT Joint Program on the Science and Policy of Global Change (E-mail: sokolov@mit.edu)

There are, however, well known problems with the use of so-called “ensembles of opportunity” (Tebaldi and Knutti 2007). Among them are difficulties with defining the relative weights of different models and the fact that existing AOGCMs do not cover the full range of uncertainty in climate sensitivity and the rate of oceanic heat uptake. By contrast, earth system models of intermediate complexity can be used effectively for producing probabilistic predictions of future climate changes, due to their computational efficiency and ability to vary the above mentioned characteristics over wide ranges.

Here we use the climate component of the MIT Integrated Global System Model (Sokolov *et al.* 2005) to evaluate the climate response and its associated uncertainty when we prescribe the forcing from four different scenarios. The probability distributions for the uncertain input parameters were obtained by comparing 20th Century temperature changes as simulated by the MIT model with available observations (Forest *et al.* 2006). In particular we use the probability distributions obtained this way when they are combined with an expert prior on climate sensitivity (Forest *et al.* 2006).

Distributions presented by Forest *et al.* (2006) are based on data for the trend in the ocean temperature averaged the 0-3km layer from Levitus *et al.* (2005). There are, however, significant differences between estimates of changes in deep ocean temperature obtained in different studies (Gouretski and Koltermann 2007; Domingues *et al.* 2008). To evaluate the impact of these differences on climate projections, we constructed two additional sets of input parameters distributions and carried out additional ensembles of 21st century climate simulations for two forcing scenarios. We also compare these projections with projections produced by the ensemble of AR4 AOGCMs.

2. MODEL DESCRIPTION

The climate model used in this study, as well as in Forest *et al.* (2006), is a modified version of the model described by Sokolov and Stone (1998). It consists of a 2-dimensional (zonally averaged) statistical-dynamical atmospheric model coupled to an ocean mixed layer model with temperature anomalies diffused below the mixed layer. The atmospheric model is derived from the Goddard Institute for Space Studies (GISS) Model II general circulation model (GCM) (Hansen *et al.* 1983) and uses parameterizations of the eddy transports of momentum, heat and moisture by baroclinic eddies (Stone and Yao 1987, 1990). The model uses the GISS radiative transfer code which contains all radiatively important trace gases as well as aerosols. The surface area of each latitude band is divided into fractions of land, ocean, land-ice and sea-ice, with the surface fluxes and surface temperature computed separately for each surface type. The version used here has 4 degree latitudinal resolution and 11 layers in the vertical. The Q-flux ocean mixed layer model and the thermodynamic sea-ice model have 4 degree by 5 degree latitude-longitude resolution and are described by Hansen *et al.* (1984).

The climate sensitivity of the atmospheric model (S) can be changed by varying the strength of the cloud feedback (Sokolov and Stone 1998), while the rate of deep oceanic heat uptake can be changed by varying the value of the global mean diffusion coefficient (K_v) used to mix ocean

temperature anomalies below the mixed layer. In spite of the simplicity of the MIT 2D model, it has been shown to be able, with the appropriate choice of the model's parameters (S , K_v) and the same forcing scenario, to reproduce the behavior of different AOGCMs over a wide range of climate sensitivities and rates of oceanic heat uptakes (Sokolov and Stone 1998; Sokolov *et al.* 2003). Fits for different AOGCMs are obtained using results from their simulations with increasing atmospheric CO_2 concentrations. Reproducing the AOGCM results for other forcing scenarios, such as the SRES scenarios, is more complicated. In some cases different sets of forcing agents were taken into account in the simulations of climate for 20th and 21st centuries with different AOGCMs. In addition, the strengths of aerosol forcing and the efficacies of different forcings vary among models.

As a result differences in the “climate forcing”¹ among AOGCMs (**Figure 1**), are significantly larger than differences in CO_2 forcing (Forster and Taylor 2006). However, if for a particular forcing scenario the “climate forcing” for a given AOGCM is similar to the “climate forcing” simulated by the MIT 2D model, then the version of the MIT model, reproducing the behavior of this AOGCM for changes in CO_2 only, also reproduces its behavior for that forcing scenarios (**Figure 2**).




Figure 1. Climate forcing for SRES A1B in the last decade of the 21st century relative to present (1980-1999) for AR4 AOGCMs, MIT IGSM and from Myhre *et al.* (2001).

¹ “Climate forcing” for a given scenario is calculated from the changes in SAT and heat flux at the top of the atmosphere using the feedback parameter estimated from the simulation with 1% per year CO_2 increase (Forster and Taylor, 2006)




Figure 2. Change in global mean annual mean surface air temperature simulated by the GFDL 2.1 model (dashed lines) and the corresponding version of the IGSM (solid lines) in the simulations with 1% per year CO₂ increase (top) and with SRES A2, A1B and B2 scenarios (bottom).

3. PROBABILITY DISTRIBUTIONS OF INPUT CLIMATE PARAMETERS

Performing probabilistic climate forecast requires knowledge of probability distributions for climate system parameters determining model response to an external radiative forcing. Forest *et al.* (2002) proposed a method for obtaining such distributions based on comparison of the results of 20th century climate simulations with observational records of surface, upper air and deep ocean temperature changes. Distributions presented by Forest *et al.* (2002, 2006 and 2008) were obtained using estimates of changes in deep ocean heat content provided by Levitus *et al.* (2005). However, several other estimates have been published in the last few years (*e.g.* Gouretski and Koltermann 2007; Ishii *et al.* 2006; Carton and Santorelli (2008); Domingues *et al.* 2008). The methods used in different studies vary in their treatment of sub-surface temperature measurements from different types of instruments (MBT, XBT and so on) as well as in the methods used to estimate changes in data-sparse regions. Only three papers, Levitus *et al.* (2005), Gouretski and Koltermann (2007) and Domingues *et al.* (2008) (LAB05, GK07 and DOM08 hereafter) provide observationally based estimates of changes in the ocean heat content for the 0-3000 m layer (**Figure 3**). All the others give results for the upper 700m only. When we re-did our analyses using trends based on just the upper 700 m we found that they did not introduce any constraints on the climate parameters beyond those resulting from using just the upper air and surface temperature data. This lack of impact is due to the strong correlation between upper-ocean and sea surface temperatures and the large natural variability of the upper ocean layers.




Figure 3. Changes in ocean heat content from LAB05 (red) , GK07 (green) and DOM08 (black) for the 0-3000m layer as estimated by each group. Five year running means were applied to the GK07 data and the errors combined appropriately. The error bars are nominally 1-sigma standard errors but represent different sources of uncertainty in each case.




Figure 4. The marginal posterior probability density function for S-Sqrt(Kv) parameter space obtained using data for surface, upper-air and deep-ocean temperatures by LAB05 (blue) and DOM08 (black) and using surface and upper-air temperatures only (No Ocean data) (red). Thick contours denote rejection regions for significance levels of 90%, 50% and 10% respectively. Red dots indicate the median values from respective 1D marginal distributions. Thin contours show values of TCR from a 250-member ensemble of simulations with the input probability distributions based on the LAB05 data.

Figure 4 shows the marginal posterior probability density functions for the S-Sqrt(Kv) parameter space ($\text{Sqrt}(Kv) = \sqrt{Kv}$) obtained using the LAB05 and DOM08 estimates for changes in deep ocean temperature and their respective estimates of the uncertainty in the trends². These estimates are respectively the smallest and largest estimates of the 0-3000m trend. We note that each of these trends lie outside the uncertainty range cited for the other trend, and thus the published uncertainty estimates are almost certainly underestimates. Therefore we also include in Figure 4 for comparison the distribution constructed without the use of any ocean data (*i.e.*, No Ocean data, hereafter, NO). We note that the probability distributions for climate parameters obtained using upper-ocean (0-700m) data are almost identical to the NO distribution. The use of the LAB05 and DOM08 analyses strongly affects the range of acceptable values of the rate of oceanic heat uptake (**Table 1**). However, due to the correlation between climate sensitivity and the rate of oceanic heat uptake imposed by the data on surface air temperature, changes in SAT in response to a 1% per year increase in CO₂ concentration in simulations with the median values of S and Kv from the different parameter distributions are not very different (see Figure 4).

Table 1. Percentiles of marginal distributions of Sqrt(Kv) obtained under different assumptions on deep ocean temperature changes.

	2.5%	5.0%	25.0%	50.0%	75.0%	95.0%	97.5%
LAB05	0.12	0.20	0.46	0.73	1.12	1.87	2.16
NO	0.14	0.24	0.75	1.47	2.45	5.39	6.44
DOM08	0.44	0.65	1.87	3.16	4.74	7.00	7.46

4. SIMULATIONS

First we carried out three 250-member ensembles of simulations from year 1860 to year 2000, using different values of the climate sensitivity, the rate of the oceanic heat uptake and the strength of the aerosol forcing for each of the distributions based on the LAB05, NO, and DOM08 results. Different combinations of model parameters controlling these characteristics were chosen using the Latin Hypercube Sampling algorithm (Iman and Helton, 1988) from the probability density functions described in the previous section. The distribution of each model parameter was divided into 250 segments of equal probability and sampling without replacement was performed, so that every segment was used once. Details on the sampling procedure can be found in Webster *et al.* (2003).

In each simulation, the MIT 2D climate model was forced by the observed changes in GHGs, stratospheric aerosols from volcanic eruptions, tropospheric and stratospheric ozone, solar irradiance and changes in vegetation due to land use change. Details of the forcings are given in Forest *et al.* (2006). As can be seen from **Figure 5**, changes in SAT in simulations with median values of climate parameters from all three distributions agree well with observations. Then we carried out a number of 250-member ensembles of simulations for several forcing scenarios

² The LAB05 analysis has recently been updated to take into account the systematic errors in the XBT data noted by GK07. The result, which is available on the NOAA website, is not appreciably different from the LAB05 result.

using different distributions of the climate parameters (**Table 2**). Each simulation was started from the end of the corresponding 20th Century simulation and simulated years 2001-2100. Data for GHG concentrations for the SRES scenarios B1, A1B and A2 scenarios were taken from the web site of the Goddard Institute for Space Studies (http://data.giss.nasa.gov/modelforce/ghgases/GCM_2004.html). In addition to changes in CO₂ (ISAM reference), N₂O, CH₄, CFC-11 and CFC-12, these data include changes in additional trace gases (Hansen and Sato, 2004). Changes in the loading of sulfate aerosols were prescribed according to the IPCC TAR's description (Houghton *et al.* 2001). Changes in black and organic carbon aerosols were not included.




Figure 5. Changes in global-mean annual-mean surface air temperature in simulations with median values of climate parameter from different distributions. Observations are from Jones (2003).

In the simulations with the MIT REF scenario we used GHG and sulfate aerosol concentrations obtained in a reference simulation with the full version of the MIT IGSM2.2 with “business as usual” emissions. Detailed information on this scenario can be found in Prinn *et al.* (2008). It is worth noting that the total radiative forcing and SAT changes in simulations with the MIT REF scenario are close to those in simulations with the SRES A1FI scenario. The chosen set of simulations allows us to explore the dependency of the projected climate changes on both the forcing scenarios and the choice of ocean data.

Table 2. List of simulations with different input parameters and different forcing scenarios.

Input parameters distributions	Forcing scenarios			
	SRES B1	SRES A1B	SRES A2	MIT REF
LAB05	X	X	X	X
NO	-	X	-	X
DOM08	-	X	-	X

5. RESULTS AND DISCUSSION

The use of different pdfs for input climate parameters has a smaller effect on the distributions of projected surface warming than on the distribution of projected sea level rise (**Figure 6, Tables 3 and 4**) because a correlation between the input parameters is imposed by the 20th century surface air temperature data. For example, the mean values of sea level rise in the ensembles with NO and DOM08 input distributions exceed the mean value for LAB05 distribution by 40 and 51%, respectively, while changes in the mean values of the increase in SAT are only 6 and 13%. The choice of input distributions also significantly changes the shape of the probability distribution for projected thermosteric sea level rise. It has been noted in several papers (*e.g.* Knutti *et al.* 2008; Meinshausen *et al.* 2008) that, for the given set of AOGCM models, the shapes of the distributions for SAT increase are similar under different SRES scenarios. Our simulations yielded similar results. Ratios of the different percentiles of the given distribution to the corresponding mean differ only slightly between ensembles with different forcing scenarios (see **Table 5**). At the same time, the ratios of the SAT changes in the simulations with different forcing scenarios for *a given version* of the MIT IGSM are not defined just by the ratios of forcings, but also depend on the values of climate sensitivity and the rate of heat uptake by the ocean.

For each of the ensembles listed in Table 2, the SAT change in a simulation using the median values of the input climate parameters is close to the median value of SAT changes from the corresponding ensemble of simulations. This fact, in combination with the similarity of the statistical properties of the distributions of projected surface warming for different forcing scenarios, allows us to approximate the probability ranges for surface warming under the SRES B1 and A2 scenarios for the NO and DOM08 sets of input parameters without running the corresponding ensembles. Namely, we run simulations with the median values of climate parameters for the remaining scenarios and then used ratios from Table 5 to estimate probability ranges. Those ranges are shown in **Figure 7** by dashed lines.

Table 3. Distributions of temperature changes in the last decade of 21st century relative to the 1981-2000 mean.

		5.0%	16.7%	50.0%	Mean	83.3%	95.0%
LAB05	B1	1.51	1.69	2.10	2.08	2.37	2.71
	A1B	2.30	2.61	3.17	3.17	3.64	4.11
	A2	3.00	3.43	4.06	4.03	4.55	4.98
	REF	3.62	4.11	4.86	4.83	5.41	6.02
NO	A1B	2.20	2.45	2.94	3.01	3.55	3.86
	REF	3.33	3.76	4.50	4.50	5.23	5.78
DOM08	A1B	2.13	2.31	2.58	2.66	3.00	3.44
	REF	3.16	3.43	3.91	3.99	4.50	5.06




Figure 6. Frequency distributions for SAT increase under SRES A1B (a) and MIT REF (b) scenarios and corresponding thermosteric sea level rise (c and d) in 2091 to 2100 relative to the 1981 to 2000 average in simulations with LAB05 (blue) DOM08 (green) and NO (red) input parameter distributions. Solid horizontal bars show 5-95% ranges from 250-member ensembles of simulations with the MIT model, dashed horizontal bars show 5-95% ranges from the multi-model IPCC AR4 AOGCM ensemble (from figure TS.27 of Solomon *et al* (2007))

Table 4. Distributions of thermosteric sea level rise in the last decade of 21st century relative year to the 1981-2000 mean.

		5.0%	16.7%	50.0%	Mean	83.3%	95.0%
LAB05	B1	5.23	8.85	13.02	12.40	17.85	21.26
	A1B	7.67	12.16	17.45	16.77	23.35	27.60
	A2	10.64	15.80	22.10	21.35	28.65	33.56
NO	REF	11.80	18.91	26.95	26.11	35.73	42.11
	A1B	11.27	14.65	24.40	24.54	33.32	38.07
	REF	17.65	23.10	37.83	37.80	51.45	58.66
DOM08	A1B	15.26	25.73	36.00	34.34	42.76	45.26
	REF	21.50	32.55	42.71	40.82	49.15	51.47

The values of Sqrt(Kv) required for the MIT IGSM to reproduce the results of different AR4 AOGCMs in simulations with 1% per year CO₂ increase, range from 0.9 cms^{-1/2} to 2.0 cms^{-1/2}. The Sqrt(Kv) values for practically all the AR4 models fall in the upper half of the Sqrt(Kv) range suggested by the pdf using the LAB05 ocean heat content data and in the low half of the range of the Sqrt(Kv) distribution based on the DOM08 results.

As a result, the simulations using the LAB05 input parameter distribution suggest stronger warming than the ensemble of the AR4 AOGCMs (see Figures 6 and 7) -- namely by the end of the 21st century (2091-2100) surface air temperature will increase above the present level (1980-1999) by 1.7C to 2.4C (16.7 to 83.3 percentiles) for B1, 2.6C to 3.6C for A1B and 3.4C to 4.6C for A2. The corresponding increases in the mean are 2.1C, 3.2C and 4.0C respectively. . From the AR4 AOGCM ensemble (also shown in Figures 6 and 7), the mean increases are 1.8C, 2.8C and 3.4C for the B1, A1B and A2 scenarios, respectively. The corresponding ranges for sea level rise due to thermal expansion of sea-water in LAB05 simulations are: 9 cm to 18 cm for B1, 12 cm to 23 cm for A1B and 16 cm to 29 cm for A2 (Table 4).

Table 5. Ratios of the percentiles values to the means for probability distributions shown in Table 3.

		5.0%	16.7%	50.0%	83.3%	95.0%
LAB05	B1	0.72	0.81	1.01	1.14	1.30
	A1B	0.73	0.82	1.00	1.15	1.30
	A2	0.74	0.85	1.01	1.13	1.23
NO	REF	0.75	0.85	1.01	1.12	1.25
	A1B	0.73	0.81	0.98	1.18	1.28
	REF	0.74	0.84	1.00	1.16	1.28
DOM08	A1B	0.80	0.87	0.97	1.13	1.29
	REF	0.79	0.86	0.98	1.13	1.27

The upper bounds of the distributions for SAT increases obtained in the LAB05 ensemble also significantly exceed the upper bounds simulated by the AR4 AOGCMs (Figure 5a). For example, the probability of surface warming exceeding 4.1 C by the end of 21st Century under

the A2 scenario is 5% according to the AR4 AOGCM ensemble (Figure TS.27 of Solomon *et al.* (2007)), but 46% according to our results. The SAT changes simulated by the AR4 AOGCMs almost completely lie below the median suggested by our LAB05 projections for the high emissions scenario (A2) (Figure 7b) and below 83.3 percentile for the other two scenarios (Figures 6a and 7a). The larger difference for A2, rather than for the other two scenarios is explained, in part, by the fact that the AR4 simulations for different SRES scenarios were carried out with different sets of AOGCMs. Thus the MIROC3.2 (hires) AOGCM, which produces the highest warming in the simulations for the A1B and B1 scenarios, was not used in the simulations with SRES A2.




Figure 7. Frequency distributions for SAT increase in 2091 to 2100 relative to the 1981 to 2000 average for the B1 (a) and A2 (b) SRES scenarios in simulations with LAB05 input parameter distribution. Solid horizontal bars show 5-95% ranges from 250-member ensembles of simulations with the MIT IGSM; green and red dashed horizontal bars show 5-95% ranges for DOM08 and NO approximated by scaling (see text for details); blue dashed horizontal bars show 5-95% ranges from the multi-model IPCC AR4 AOGCM ensemble (from figure TS.27 of Solomon *et al.*, 2007)

On the other hand our projections using the DOM08 analysis are more consistent with the AR4 AOGCM projections. However our NO projections are closer to our LAB05-based projections, which indicates that, for the assumed prior for climate sensitivity, the observed changes in surface and upper air temperature are more consistent with the weaker ocean warming trend in the LAB05 analysis than with the stronger trend in the DOM08 analysis.

The differences in forcing between AOGCMs (illustrated in Figure 1) are unlikely to affect our comparison in a significant way for two reasons. First, the climate forcing in the MIT IGSM simulations with mean values of climate sensitivity, rate of oceanic heat uptake and strength of aerosol forcing turned out to be very similar to the forcing averaged over the set of AR4 AOGCMs (see Figure 1). Second, the correlation between TCR, which provides a good measure of model response to an external forcing on time scales of 50-100 years, and the strength of climate forcing for the AR4 AOGCMs is very weak.

As noted above, the use of input distributions based on different estimates of changes in the ocean heat content has a strong effect on the projected sea level rise due to thermal expansion. The range of values produced by the AR4 AOGCMs for the SRES A1B scenario falls in between ranges obtained in the different MIT ensembles (Figure 6c).

6. CONCLUSIONS

Uncertainties in the estimates of the 20th century changes in the deep ocean heat content have a strong effect on projections of the thermosteric sea level rise. The median value of the thermosteric sea level rise under the MIT reference forcing scenario is 41 cm if the DOM08 data is used, but 27 cm if the LAB05 data is used. The effect on SAT increases is more modest because of the correlation between climate sensitivity and ocean heat uptake required by observed 20th century temperature changes. Nevertheless the effect is still significant. The median SAT increase at the end of the 21st century for the MIT reference scenario is 3.9 C if the DOM08 data are used for producing the probability distributions, but 4.9 C if the LAB05 data are used.

For the forcing scenarios used in this study, the statistical properties of the distributions of changes in surface air temperature do not depend on the choice of distribution for input climate parameters (see Table 5). Such similarity in the distributions for projected surface warming is explained by the correlation between S and $\text{Sqrt}(Kv)$ imposed by the transient changes in SAT observed during 20th century and by the fact that in all scenarios considered here forcing increases monotonically with time. Distributions for the equilibrium SAT changes under scenarios where the forcing is stabilized will be defined primarily by the input distributions for climate sensitivity.

The estimates of SAT increase produced by the multi-model ensemble of the IPCC AR4 AOGCMs slightly overestimate the results of the MIT ensemble for input distributions based on the DOM08 data, and strongly underestimate the surface warming suggested by the MIT results using the LAB05 data. For the sea level rise due to thermal expansion the situation is the opposite. As noted above, the LAB05 analysis is more consistent with the observed 20th century surface and upper air temperature changes. These results emphasize the necessity for producing reliable estimates for changes of deep ocean heat content.

Acknowledgments

We would like to thank Piers Forster and Karl Taylor for providing data on climate forcing for AR4 AOGCMs, Catia Domingues for providing data on ocean heat content, and Sharmila Ganguly for help with producing this report. This work was supported in part by the Office of Science (BER), U.S. Dept. of Energy Grant No. DE-FG02-93ER61677, NSF, and by the MIT Joint Program on the Science and Policy of Global Change. We thank Marcus Sarofim for LHS samples of input parameters and Josh Willis for discussions on the ocean heat content data. We acknowledge the modeling groups, the Program for Climate Model Diagnosis and Intercomparison (PCMDI) and the WCRP's Working Group on Coupled Modelling (WGCM) for their roles in making available the WCRP CMIP3 multi-model dataset. Support of this dataset is provided by the Office of Science, U.S. Department of Energy

7. REFERENCES

- Andronova, N. and M.E. Schlesinger, (2001): Objective estimation of the probability distribution for climate sensitivity. *J. Geophys. Res.* **106 (D19)**: 22605-22612, [doi:10.1029/2000JD000259](https://doi.org/10.1029/2000JD000259)
- Carton, J.A. and A. Santorelli, (2008): Global Decadal Upper Ocean Heat Content as Viewed in Nine Analyses. *J. Climate* in press http://ams.allenpress.com/archive/1520-0442/preprint/2008/pdf/10.1175_2008JCLI2489.1.pdf
- Domingues, C.M., J.A. Church, N.J. White, P.J. Gleckler, S.E. Wijffels, P.M. Barker, and Jeff R. Dunn, (2008) Improved estimates of upper-ocean warming and multi-decadal sea-level rise. *Nature*, **453**: 1090-1094.
- Forest, C. E., P.H. Stone, A.P. Sokolov, M.R. Allen and M. Webster, (2002): Quantifying uncertainties in climate system properties with the use of recent climate observations. *Science*, **295**: 113-117.
- Forest, C.E., P.H. Stone and A.P. Sokolov, (2006): Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings. *Geophys. Res. Lett.*, **33**, L01705, <http://dx.doi.org/doi:10.1029/2005GL023977>
- Forest, C. E., P.H. Stone and A.P. Sokolov, (2008): Constraining climate model parameters from observed 20th century changes, *Tellus*, **60A**: 911-920.
- Forster, P.M.D. and K.E. Taylor, (2006) Climate forcing and climate sensitivities diagnosed from coupled climate model integrations, *J. Clim.*, **19**: 6181-6194.
- Frame, D.J., B.B.B. Booth, J.A. Kettleborough, D.A. Stainforth, J.M. Gregory, M. Collins and M.R. Allen, (2005): Constraining climate forecasts: the role of prior assumptions. *Geophys. Res. Lett.* **32**, L09702. [doi:10.1029/2004GL022241](https://doi.org/10.1029/2004GL022241)
- Gouretski, V., and K.P. Koltermann, (2007): How much is the ocean really warming? *Geophys. Res. Lett.*, **34**, L01610, <http://dx.doi.org/doi:10.1029/2006GL027834>
- Hansen, J., G. Russell, D. Rind, P. Stone, A. Lacis, S. Lebedeff, R. Ruedy and L. Travis, (1983): Efficient three-dimensional global models for climate studies: Models I and II. *Mon. Wea. Rev.*, **111**: 609–662.
- Hansen, J, A. Lacis, D. Rind, G. Russell, P. Stone, I. Fung, R. Ruedy and J. Lerner, (1984): Climate sensitivity: Analysis of feedback mechanisms. In *Climate Processes and Climate Sensitivity*, *Geophys. Monogr.*, **No. 29**, Amer. Geophys. Union: 130–163.
- Hansen, J., and M. Sato, (2004): [Greenhouse gas growth rates](https://doi.org/10.1073/pnas.0406024101). *Proc. Natl. Acad. Sci. USA* **101**: 16109-16114.
- Houghton, Y. Ding, D.J. Griggs, M. Noguer, P. J. van der Linden and D. Xiaosu (Eds.) (2001): *Climate Change 2001: The Scientific Basis*. Cambridge University Press, UK. pp 944.
- Iman, R.L. and J.C. Helton, (1988): An investigation of uncertainty and sensitivity analysis techniques for computer models, *Risk Anal.*, **8**: 71-90.
- Jones, P.D., and A. Moberg, (2003): Hemispheric and Large-Scale Surface Air Temperature Variations: An Extensive Revision and an Update to 2001, *J. Climate*, **16**, 206-223.
- Knutti, R., G.A. Meehl, M.R. Allen and D.A. Stainforth, (2006): Constraining climate sensitivity from the seasonal cycle in surface temperature. *J. Clim.* **19**: 4224–4233.
- Levitus, S., J. Antonov and T.P. Boyer, (2005): Warming of the World Ocean 1955–2003. *Geophys. Res. Lett.*, **32**, L02604, <http://dx.doi.org/doi:10.1029/2004GL021592>
- Meehl, G.A., C. Covey, T. Delworth, M. Latif, B. McAvaney, J.F.B. Mitchell, R.J. Stouffer and K.E. Taylor, (2007a): The WCRP CMIP3 Multimodel Dataset: A New Era in Climate Change Research. *Bull. American Meteorological Society*, **88**(9), 1383–1394.

- Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao, (2007b): Global Climate Projections. In: Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Meinshausen, M., S.C.B. Raper and T.M.L. Wigley, (2008): Emulating IPCC AR4 atmosphere-ocean and carbon cycle models for projecting global-mean, hemispheric and land/ocean temperatures: MAGICC 6.0. *Atmospheric Chemistry and Physics Discussions*, **8**: 6153-6272, <http://www.atmos-chem-phys-discuss.net/8/6153/2008/acpd-8-6153-2008.html>
- Prinn, R, S. Paltsev, A. Sokolov, M. Sarofim, J. Reilly, and H. Jacoby, (2008): The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model, MIT JP Report 163. http://globalchange.mit.edu/files/document/MITJPSPGC_Rpt163.pdf
- Pierce, D.W., T.P. Barnett, K.M. AchutaRao, P.J. Gleckler, J.M. Gregory, and W.M. Washington, (2006): Anthropogenic Warming of the Oceans: Observations and Model Results. *J. Clim.*, **19**: 1873–1900.
- Randall, D.A. *et al.* (2007): Climate Models and Their Evaluation. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., M. Tignor, & Miller, H.L. (eds.), *Climate Change 2007, The Physical Science Basis*. Cambridge University Press, Cambridge, UK.
- Sokolov, A.P. and P.H. Stone, (1998): A flexible climate model for use in integrated assessments. *Clim. Dyn.*, **14**: 291–303.
- Sokolov, A., C.E. Forest and P.H. Stone, (2003): Comparing oceanic heat uptake in AOGCM transient climate change experiments. *J. Clim.*, **16**: 1573-1582.
- Sokolov, A.P. *et al.* (2005): The MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation, MIT JP Report 124. http://globalchange.mit.edu/files/document/MITJPSPGC_Rpt124.pdf
- Solomon S. *et al.* (2007) Technical Summary. In: *Climate Change (2007): The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Stone, P. H. and M.-S. Yao, (1987): Development of a two-dimensional zonally averaged statistical–dynamical model. Part II: The role of eddy momentum fluxes in the general circulation and their parameterization. *J. Atmos. Sci.*, **44**: 3769–3786.
- Stone, P.H. and M.-S. Yao, (1990): Development of a two-dimensional zonally averaged statistical–dynamical model. Part III: The parameterization of the eddy fluxes of heat and moisture. *J. Clim.*, **3**: 726–740.
- Tebaldi, C. and R. Knutti, (2007): The use of the multi-model ensemble in probabilistic climate projections. *Phil. Trans. R. Soc. A* **365**: 2053-2075 <http://dx.doi.org/doi:10.1098/rsta.2007.2076>
- Webster, M. *et al.* (2003): Uncertainty analysis of climate change and policy response, *Clim. Change*, **61**: 295-320.
- Wijffels, S.E. *et al.* (2008): Changing expendable Bathythermograph Fallrates and their Impact on Estimates of Thermosteric Sea Level Rise. *J. Climat.* **21**: 5657–5672

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

1. **Uncertainty in Climate Change Policy Analysis**
Jacoby & Prinn December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO₂ Concentration** *Xiao et al.* October 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* January 1996
5. **World Energy Consumption and CO₂ Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996 (*superseded* by No. 125)
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 124)
8. **Relative Roles of Changes in CO₂ and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO₂ Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N₂O and CH₄ from the Terrestrial Biosphere to the Atmosphere** *Liu* Aug. 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes** *Xiao et al.* November 1996
13. **Greenhouse Policy Architectures and Institutions** *Schmalensee* November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO₂ Capture and Disposal** *Eckaus et al.* December 1996
16. **What Drives Deforestation in the Brazilian Amazon?** *Pfaff* December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Subgrid Scale Processes in Global Atm. Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming** *Jacoby, Prinn & Schmalensee* August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake and their Impact on Climate Projections** *Sokolov et al.* September 1997
24. **A Global Interactive Chemistry and Climate Model** *Wang, Prinn & Sokolov* September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry & Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements** *Yang & Jacoby* October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere** *Xiao et al.* November 1997
29. **Analysis of CO₂ Emissions from Fossil Fuel in Korea: 1961-1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby et al.* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions and Resultant Climatic Changes on Atmospheric OH** *Wang & Prinn* April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies** *Prinn et al.* June 1998
37. **Quantifying the Uncertainty in Climate Predictions** *Webster & Sokolov* July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* September 1998
39. **Uncertainty in Atmospheric CO₂ (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO₂ Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* Oct. 1998
41. **The Effects on Developing Countries of the Kyoto Protocol and CO₂ Emissions Trading** *Ellerman et al.* November 1998
42. **Obstacles to Global CO₂ Trading: A Familiar Problem** *Ellerman* November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* November 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol** *Reilly et al.* January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques** *Forest et al.* April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture** *Eckaus & Tso* July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement** *Babiker, Reilly & Ellerman* August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999
53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** *Babiker & Jacoby* October 1999

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** *Bugnion* Oct 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** *Bugnion* October 1999
56. **The Kyoto Protocol and Developing Countries** *Babiker et al.* October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** *Bugnion & Reiner* November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** *Reilly, Mayer & Harnisch* March 2000
59. **Supplementarity: An Invitation for Monopsony?** *Ellerman & Sue Wing* April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** *Kamenkovich et al.* May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** *Babiker et al.* May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** *Forest et al.* May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** *Mayer et al.* June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** *Lahiri et al.* Aug 2000
65. **Rethinking the Kyoto Emissions Targets** *Babiker & Eckaus* August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** *Viguié* September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** *Webster* October 2000
68. **How to Think About Human Influence on Climate** *Forest, Stone & Jacoby* October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** *Ellerman* Nov 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** *Viguié et al.* February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** *Babiker et al.* February 2001 (*superseded* by No. 125)
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** *Fullerton & Metcalf* March '01
73. **Uncertainty Analysis of Global Climate Change Projections** *Webster et al.* Mar. '01 (*superseded* by No. 95)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** *Babiker et al.* June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO₂** *Kamenkovich et al.* July 2001
76. **CO₂ Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** *Ellerman & Tsukada* July 2001
77. **Comparing Greenhouse Gases** *Reilly et al.* July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** *Forest et al.* July 2001
79. **Uncertainty in Emissions Projections for Climate Models** *Webster et al.* August 2001
80. **Uncertainty in Atmospheric CO₂ Predictions from a Global Ocean Carbon Cycle Model** *Holian et al.* September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** *Sokolov et al.* December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** *Babiker, Jacoby & Reiner* February 2002
83. **The "Safety Valve" and Climate Policy** *Jacoby & Ellerman* February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** *Wang* March 2002
85. **Tax Distortions and Global Climate Policy** *Babiker et al.* May 2002
86. **Incentive-based Approaches for Mitigating Greenhouse Gas Emissions: Issues and Prospects for India** *Gupta* June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** *Huang, Stone & Hill* September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** *Huang et al.* September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Information** *McFarland et al.* October 2002
90. **Ozone Effects on Net Primary Production and Carbon Sequestration in the U.S. Using a Biogeochemistry Model** *Felzer et al.* November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** *Carlén* November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** *Herzog et al.* December 2002
93. **Is International Emissions Trading Always Beneficial?** *Babiker et al.* December 2002
94. **Modeling Non-CO₂ Greenhouse Gas Abatement** *Hyman et al.* December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** *Webster et al.* December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** *Carlén* January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the United States: The McCain-Lieberman Proposal** *Paltsev et al.* June 2003
98. **Russia's Role in the Kyoto Protocol** *Bernard et al.* Jun '03
99. **Thermohaline Circulation Stability: A Box Model Study** *Lucarini & Stone* June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** *Ellerman & Sue Wing* July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** *Schafer & Jacoby* July 2003
102. **Induced Technical Change and the Cost of Climate Policy** *Sue Wing* September 2003
103. **Past and Future Effects of Ozone on Net Primary Production and Carbon Sequestration Using a Global Biogeochemical Model** *Felzer et al.* (revised) January 2004
104. **A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** *Zhuang et al.* November 2003

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

105. **Analysis of Strategies of Companies under Carbon Constraint** *Hashimoto* January 2004
106. **Climate Prediction: The Limits of Ocean Models** *Stone* February 2004
107. **Informing Climate Policy Given Incommensurable Benefits Estimates** *Jacoby* February 2004
108. **Methane Fluxes Between Terrestrial Ecosystems and the Atmosphere at High Latitudes During the Past Century** *Zhuang et al.* March 2004
109. **Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** *Dalan et al.* May 2004
110. **Stabilization and Global Climate Policy** *Sarofim et al.* July 2004
111. **Technology and Technical Change in the MIT EPPA Model** *Jacoby et al.* July 2004
112. **The Cost of Kyoto Protocol Targets: The Case of Japan** *Paltsev et al.* July 2004
113. **Economic Benefits of Air Pollution Regulation in the USA: An Integrated Approach** *Yang et al.* (revised) Jan. 2005
114. **The Role of Non-CO₂ Greenhouse Gases in Climate Policy: Analysis Using the MIT IGSM** *Reilly et al.* Aug. '04
115. **Future U.S. Energy Security Concerns** *Deutch* Sep. '04
116. **Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** *Sue Wing* Sept. 2004
117. **Modeling the Transport Sector: The Role of Existing Fuel Taxes in Climate Policy** *Paltsev et al.* November 2004
118. **Effects of Air Pollution Control on Climate** *Prinn et al.* January 2005
119. **Does Model Sensitivity to Changes in CO₂ Provide a Measure of Sensitivity to the Forcing of Different Nature?** *Sokolov* March 2005
120. **What Should the Government Do To Encourage Technical Change in the Energy Sector?** *Deutch* May '05
121. **Climate Change Taxes and Energy Efficiency in Japan** *Kasahara et al.* May 2005
122. **A 3D Ocean-Seaice-Carbon Cycle Model and its Coupling to a 2D Atmospheric Model: Uses in Climate Change Studies** *Dutkiewicz et al.* (revised) November 2005
123. **Simulating the Spatial Distribution of Population and Emissions to 2100** *Asadoorian* May 2005
124. **MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation** *Sokolov et al.* July 2005
125. **The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4** *Paltsev et al.* August 2005
126. **Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings** *Forest et al.* September 2005
127. **An Analysis of the European Emission Trading Scheme** *Reilly & Paltsev* October 2005
128. **Evaluating the Use of Ocean Models of Different Complexity in Climate Change Studies** *Sokolov et al.* November 2005
129. **Future Carbon Regulations and Current Investments in Alternative Coal-Fired Power Plant Designs** *Sekar et al.* December 2005
130. **Absolute vs. Intensity Limits for CO₂ Emission Control: Performance Under Uncertainty** *Sue Wing et al.* January 2006
131. **The Economic Impacts of Climate Change: Evidence from Agricultural Profits and Random Fluctuations in Weather** *Deschenes & Greenstone* January 2006
132. **The Value of Emissions Trading** *Webster et al.* Feb. 2006
133. **Estimating Probability Distributions from Complex Models with Bifurcations: The Case of Ocean Circulation Collapse** *Webster et al.* March 2006
134. **Directed Technical Change and Climate Policy** *Otto et al.* April 2006
135. **Modeling Climate Feedbacks to Energy Demand: The Case of China** *Asadoorian et al.* June 2006
136. **Bringing Transportation into a Cap-and-Trade Regime** *Ellerman, Jacoby & Zimmerman* June 2006
137. **Unemployment Effects of Climate Policy** *Babiker & Eckaus* July 2006
138. **Energy Conservation in the United States: Understanding its Role in Climate Policy** *Metcalfe* Aug. '06
139. **Directed Technical Change and the Adoption of CO₂ Abatement Technology: The Case of CO₂ Capture and Storage** *Otto & Reilly* August 2006
140. **The Allocation of European Union Allowances: Lessons, Unifying Themes and General Principles** *Buchner et al.* October 2006
141. **Over-Allocation or Abatement? A preliminary analysis of the EU ETS based on the 2006 emissions data** *Ellerman & Buchner* December 2006
142. **Federal Tax Policy Towards Energy** *Metcalfe* Jan. 2007
143. **Technical Change, Investment and Energy Intensity** *Kratena* March 2007
144. **Heavier Crude, Changing Demand for Petroleum Fuels, Regional Climate Policy, and the Location of Upgrading Capacity** *Reilly et al.* April 2007
145. **Biomass Energy and Competition for Land** *Reilly & Paltsev* April 2007
146. **Assessment of U.S. Cap-and-Trade Proposals** *Paltsev et al.* April 2007
147. **A Global Land System Framework for Integrated Climate-Change Assessments** *Schlosser et al.* May 2007
148. **Relative Roles of Climate Sensitivity and Forcing in Defining the Ocean Circulation Response to Climate Change** *Scott et al.* May 2007
149. **Global Economic Effects of Changes in Crops, Pasture, and Forests due to Changing Climate, CO₂ and Ozone** *Reilly et al.* May 2007
150. **U.S. GHG Cap-and-Trade Proposals: Application of a Forward-Looking Computable General Equilibrium Model** *Gurgel et al.* June 2007
151. **Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks between Climate and the Terrestrial Carbon Cycle** *Sokolov et al.* June 2007
152. **Energy Scenarios for East Asia: 2005-2025** *Paltsev & Reilly* July 2007
153. **Climate Change, Mortality, and Adaptation: Evidence from Annual Fluctuations in Weather in the U.S.** *Deschênes & Greenstone* August 2007

REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

- 154. Modeling the Prospects for Hydrogen Powered Transportation Through 2100** *Sandoval et al.*
February 2008
- 155. Potential Land Use Implications of a Global Biofuels Industry** *Gurgel et al.* March 2008
- 156. Estimating the Economic Cost of Sea-Level Rise**
Sugiyama et al. April 2008
- 157. Constraining Climate Model Parameters from Observed 20th Century Changes** *Forest et al.* April 2008
- 158. Analysis of the Coal Sector under Carbon Constraints** *McFarland et al.* April 2008
- 159. Impact of Sulfur and Carbonaceous Emissions from International Shipping on Aerosol Distributions and Direct Radiative Forcing** *Wang & Kim* April 2008
- 160. Analysis of U.S. Greenhouse Gas Tax Proposals**
Metcalf et al. April 2008
- 161. A Forward Looking Version of the MIT Emissions Prediction and Policy Analysis (EPPA) Model**
Babiker et al. May 2008
- 162. The European Carbon Market in Action: Lessons from the first trading period** Interim Report
Convery, Ellerman, & de Perthuis June 2008
- 163. The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model** *Prinn et al.*
September 2008
- 164. Marginal Abatement Costs and Marginal Welfare Costs for Greenhouse Gas Emissions Reductions: Results from the EPPA Model** *Holak et al.* November 2008
- 165. Uncertainty in Greenhouse Emissions and Costs of Atmospheric Stabilization** *Webster et al.* November 2008
- 166. Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content** *Sokolov et al.* November 2008